WORKSHEET 1 Summary Score Sheet

SITE INFORMATION:

Name: Orchard Middle School Address: 1024 Orchard Avenue

City: Wenatchee County: Chelan State: WA Zip: 98801

Section/Township/Range: S4/T22N/R20E

Latitude: 47° 25' 37" N Longitude: 120° 19' 35" W

TCP ID #: 771247

Site scored/ranked for the February 2006 update.

SITE DESCRIPTION (management areas, substances of concern, and quantities):

Orchard Middle School is located in a residential area in the center of Wenatchee, WA. The Middle School hosts approximately 500 sixth-, seventh-, and eighth-grade students,

This site was included in an area-wide lead and arsenic sampling program which involved collecting samples from schools suspected of having a history of past pesticide use. Prior to the mid-1940s, lead arsenate was the most widely used chemical used to control cottling moths on fruit trees. Lead (Pb) and arsenic (As) are known to be very stable in soil and tend to stay near the surface. Because of this historical background, it was suspected that the soil in the school playground might be contaminated with Pb and As. The Washington Department of Ecology (Ecology) obtained permission from Wenatchee School District to sample and test the soils for lead and arsenic from all of the Orchard Middle School grounds.

The soils throughout the property were sampled by the Department of Ecology between July and August, 2002. Samples were taken from the top six inches using a core sampler. The samples were analyzed for lead and arsenic using X-Ray Fluorescence Spectroscopy.

Sampling results at Orchard Middle School indicate that contaminant levels in soil exceed the Model Toxics Control Act Method A cleanup levels for lead (250 ppm) and/or arsenic (20 ppm) in 12 of 15 soil samples. The highest levels of arsenic and lead detected at the site were 90.5 ppm and 330 ppm, respectively. These concentrations require the site be scored and ranked under the Washington Ranking Method (WARM).

SPECIAL CONSIDERATIONS (include limitations in site file data or data which cannot be accommodated in the model, but which are important in evaluating the risk associated with the site, or any other factor(s) over-riding a decision of no further action for the site):

Due to the nature of metals in soil, it is assumed that lead and arsenic are likely not present at high concentrations at depths greater than four feet. This has been documented at several sites and seems

consistent for sites where lead arsenate pesticide was used. Samples were collected at a depth of six inches; however, for the purposes of this assessment, it is assumed that contamination extends to a depth of three feet.

Because of the presence of children at the school grounds, ingestion of contaminated soil is of concern and is considered the greatest risk to children. WARM scoring does not consider ingestion as an exposure route. However, the use of school grounds by children and their incidental ingestion of lead and arsenic contaminated soil are being addressed by Ecology through remediation efforts and by the State Department of Health through education.

ROUTE SCORES:

Surface Water/Human Health:	9.9	Surface Water/Environmental.:	13.3
Air/Human Health:	3.9	Air/Environmental:	NS
Groundwater/Human Health:	55.6		

OVERALL RANK: 3

WORKSHEET 2 Route Documentation

1. SURFACE WATER ROUTE

a. List those substances to be <u>considered</u> for scoring:

Source: 1

Lead and arsenic

b. Explain basis for choice of substance(s) to be <u>used</u> in scoring.

Analytical results from soil sampling indicate the presence of these hazardous substances at levels which exceed our current Method A cleanup levels.

c. List those management units to be considered for scoring:

Source 1

Surface and subsurface soils

d. Explain basis for choice of unit to be used in scoring:

Spills/discharges caused soil contamination

2. AIR ROUTE

a. List those substances to be considered for scoring:

Source: 1

Lead and arsenic

b. Explain basis for choice of substance(s) to be used in scoring:

Analytical results from soil sampling indicate the presence of these hazardous substances at levels which exceed our current Method A cleanup levels.

c. List those management units to be <u>considered</u> for scoring:

Source: 1

Surface and subsurface soils

d. Explain basis for choice of unit to be used in scoring:

Spills/discharges caused soil contamination

3. GROUNDWATER ROUTE

a. List those substances to be <u>considered</u> for scoring:

Source: 1

Lead and arsenic

b. Explain basis for choice of substance(s) to be used in scoring:

Analytical results from soil sampling indicate the presence of these hazardous substances at levels which exceed our current Method A cleanup levels.

c. List those management units to be <u>considered</u> for scoring:

Source: 1

Surface and subsurface soils

d. Explain basis for choice of unit to be used in scoring:

Spills/discharges caused soil contamination

WORKSHEET 4 Surface Water Route

1.0 SUBSTANCE CHARACTERISTICS

		Drinking		Acute		Chronic		Carcino	genicity	
	Substance	Water Standard (µg/L)	Value	Toxicity (mg/kg-bw)	Value	Toxicity (mg/kg/day)	Value	WOE	PF*	Value
1	Lead	5	8	NA	ND	0.001	10	ND	ND	ND
2	Arsenic	10	8	763 (rat)	5	0.001	5	1.0	1.75	7

*Potency Factor

Source: <u>1, 2</u>

Highest Value: 10 (Max = 10)

Plus 2 Bonus Points? 2 Final Toxicity Value: 12 (Max = 12)

1.2	Environmental Toxicity				
. Substance		Acute Water Quality Criteria		Mamma	Human lian Acute xicity
		(μg/L)	- Value	(mg/kg)	Value
1	Lead	82	6	ND	
2	Arsenic	360	4	ND	

Source: <u>1, 2</u>

Highest Value: $\underline{6}$ (Max = 10)

1.3 Substance Quantity (areal extent)	
Explain Basis: Fifteen samples were collected throughout the site. Each sample represented approximately 5600 square feet. Twelve sample exceeded MTCA cleanup levels. Therefore, by calculating 12 * 5600 sq. ft, 67200 sq. feet can be estimated as contaminated.	Source: 3, 8 Value: 7 (Max = 10)

2.0 MIGRATION POTENTIAL

		Source	Value
	Containment: Management unit scored as a spills/discharges/contaminated soil at the surface, with ineffectively maintained run-on/runoff controls (vegetated buffer).		4
2.1	Explain basis: While a portion of the site is capped (parking lots, buildings, etc), soil samples were collected from only uncapped areas (i.e. in bare soil or beneath grass) at a depth of six inches.	1,3	(Max = 10)
2.2	Surface Soil Permeability: the site consists of burch loam	3, 10	3 (Max = 7)
2.3	Total Annual Precipitation: The average annual precipitation for Wenatchee, WA is 8.85 in	9	1 (Max = 5)
2.4	Max 2yr/24hr Precipitation: 1.5 inches	3	2 (Max = 5)
2.5	Flood Plain: in 500-year flood plain	3, 11	1 (Max = 2)
2.6	Terrain Slope: (755' -620')/4500' = 3%	3, 10	2 (Max = 5)

3.0 TARGETS

		Source	Value
3.1	Distance to Surface Water: ~4500 feet	3, 6	4 (Max = 10)
3.2	Population Served within 2 miles (see WARM Scoring Manual Regarding Direction): $\sqrt{18} = 4.2$	3,6	4 (Max = 75)
3,3	Area Irrigated by surface water within 2 miles : $(0.75)*\sqrt{\text{# acres}} = 0.75*\sqrt{24} = 3.7$	3,6	4 (Max = 30)
3.4	Distance to Nearest Fishery Resource: ~4500 feet	3,6	6 (Max = 12)
3.5	Distance to, and Name(s) of, Nearest Sensitive Environment(s): ~4500 ft	3,6	6 (Max = 12)

4.0 RELEASE

Explain Basis: Not documented	, , , , , , , , , , , , , , , , , , ,	 	Source: <u>1, 3</u>
			Value: $\underline{0}$ (Max = $\underline{5}$)

WORKSHEET 5 Air Route

1.0 SUBSTANCE CHARACTERISTICS

Introduction 1.1.

, , , , , , , , , , , , , , , , , , ,	Air	Standard Value T		Acute Toxicity Value (mg/ m³)			Carcinogenicity		** Y
Substance	Standard (µg/m³)					Value	WOE	PF*	Value
1 Lead	0.5	10	ND		0.001	10	B2	ND	
2 Arsenic	0.00023	10	ND		0.001	5	Α	50	9

^{*} Potency Factor

Source: <u>1, 2, 3</u>

Highest Value: 10

 $(Max = \overline{10})$

Plus 2 Bonus Points? 2

Final Toxicity Value: 12 (Max = 12)

1.3 Mobility (Use numbers to refe	er to above listed subs	tances)	
1.3.1 Gaseous Mobility	1	.3.2 Particulate Mobility	
Vapor Pressure(s) (mmHg)	Soil Type	Erodibility	Climatic Factor
1	Burch loam	56	10 - 30

Source: NA Value: NS (Max = 4) Source: 10 Value: 1 $(Max = \overline{4})$

1.4 Highest Human Health Toxicity/ Mobility Matrix Value (from Table A-7)

Final Matrix Value: 6

 $(Max = 2\overline{4})$

1.5	Environmental Toxicity/Mobility					
	Substance	Non-human Mammalian Inhalation Toxicity (mg/m³)	Acute Value	Mobility (mmHg)	Value	Matrix Value
1	Lead	ND		ND		
2	Arsenic	ND		ND		

Highest Environmental Toxicity/Mobility Matrix Value (from Table A-7) = Final Matrix Value: NS (Max = 24)

1.6 Substance Quantity (areal extent)	
Explain Basis: ~67200 sq ft (see Surface Water Route 1.3)	Source: <u>3, 8</u>
	Value: 7
	(Max = 10)

2.0 MIGRATION POTENTIAL

<u> </u>		Source	Value
2.1 Containment:	Uncontaminated soil cover <2 feet thick	8	5 (Max = 10)

3.0 TARGETS

<u> </u>		Source	Value_
3.1	Nearest Population: <1000 feet to school	3, 7	10 (Max = 10)
520 (\$200)	Distance to [and name(s) of] nearest sensitive environment(s) [fisheries excluded]: NA since not scoring environmental route		NA (Max = 7)
3.3	Population served within 0.5 miles: $\sqrt{\text{pop.}} = 18 \text{ homes } * 3 = 54 \text{ people}$ $\sqrt{54} = 7.3$	3, 6	7 (Max = 75)

4.0 RELEASE

Explain Basis for scoring a release to air: Not documented	Source: <u>3, 7</u>
	Value: <u>0</u>
	(Max = 5)

WORKSHEET 6 Groundwater Route

1.0 SUBSTANCE CHARACTERISTICS

1.2	1.2 Human Toxicity									
		Drinking Acute Chronic Carcin		Carcino	ogenicity					
Substance		Water Standard (μg/L)	Value	Toxicity (mg/ kg-bw)	Value	Toxicity (mg/kg/day)	Value	WOE	PF*	Value
1	Lead	5	8	NA	ND	0.001	10	ND	ND	ND
2	Arsenic	10	8	763 (rat)	5	0.001	5	1.0	1.75	7

^{*} Potency Factor

Source: 1, 2

Highest Value: 10

 $(Max = \overline{10})$

Plus 2 Bonus Points? 2

Final Toxicity Value: 12

(Max = 12)

1.2 Mobility (use numbers to refer to above lis	ted substances)
Cations/Anions [Coefficient of Aqueous Migration (K)] Ol	R Solubility (mg/L)
1= K > 1.0 = 2	1=
2= K is 0.1 to 1.0 = 3	2=

Source: $\underline{3}$ Value: $\underline{3}$

(Max = 3)

1.3 Substance Quantity (volume):

Explain basis: Fifteen samples were collected throughout the site. Each sample represented approximately 5600 square feet. Twelve sample exceeded MTCA cleanup levels. Therefore, by calculating 12 * 5600 sq. ft, 67200 sq. feet can be estimated as contaminated. Based on previous sampling sessions where lead/arsenic has been detected to depths up to 4', Ecology has determined that is likely that contaminated soil extends to a depth of 3' bgs. Therefore the estimated volume of contamination is calculated as follows.

Source: <u>1, 3</u> **Value: <u>5</u>**

(Max=10)

67200' * 3' = 201600 ft³ or \sim 7467 yd³

2.0 MIGRATION POTENTIAL

		Source	Value
2.1	Containment (explain basis): While a portion of the site is capped (parking lots, buildings, etc), soil samples were collected from only uncapped areas (i.e. in bare soil or beneath grass) at a depth of six inches.	3	10 (Max = 10)
2.2	Net precipitation: $5.7" - 2.7" = 3.0"$	4	1 (Max = 5)
2.3	Subsurface hydraulic conductivity: the site consists of burch loam	-3	3 (Max = 4)
2.4	Vertical depth to groundwater: the nearest water wells report static groundwater levels at 12 and 36 feet below ground surface	3,5	6 (Max = 8)

3.0 TARGETS

		Source	Value
3.1	Groundwater usage: Public supply, but alternate sources available with minimum hookup requirements	3	4 (Max = 10)
3.2	Distance to nearest drinking water well: ~3700 feet	3, 5	2 (Max = 5)
3,3	Population served within 2 miles: √ pop. = <10,000	3, 7	100 (Max = 100)
3.4	Area irrigated by (groundwater) wells within 2 miles: $(0.75)*\sqrt{\# \text{ acres}} = \underbrace{0.75*\sqrt{3}=1.3}$	3, 6	1 (Max = 50)

4.0 RELEASE

Source	Value
	0 (Max = 5)
	Source

SOURCES USED IN SCORING

- 1. Analytical results of soil sampling conducted between July and August 2002 by the WA State Dept. of Ecology
- 2. Washington State Department of Ecology, Toxicology Database for Use in Washington Ranking Method Scoring, January 1992
- 3. Washington State Department of Ecology, WARM Scoring Manual, April 1992.
- 4. Washington Climate Net Rainfall Table
- 5. Washington State Department of Ecology, Water Well Reports
- 6. Washington State Department of Ecology, Water Rights Application System (WRATS) printouts
- 7. Washington State Department of Health, Office of Drinking Water Sentry website printout for public water supplies
- 8. Orchard Middle School file, WSDOE records at the Central Regional Office

- 9. Western Regional Climate Center's Historical Climate Information
- 10. National Resources Conservation Service Soil Survey of Chelan County Area, WA
- 11. FEMA online flood data